Bio 1M03 Version Q

1. Which of the following would be the *least* likely trigger for an adaptive radiation in a given clade of lizards?

- **A.** They arrive on an island that has no ecologically similar species
- B. Small snakes that compete with the lizards arrive on their island
- C. Large snakes that eat the lizards arrive on their island
- **D.** A new mutation in the lizards that opens the way for further mutations
- **E.** A new mutation in the plants on their island that opens the way for further mutations
- **2.** A certain valley has more fossils of snails than of slugs. Which of the following is *not* a likely reason for this pattern?
 - A. The valley had more snails than slugs over the relevant period of time
 - **B.** The snails fossilize better because of their hard shells
 - C. The snails lived in a wetter part of the valley
 - **D.** The snails spent less time underground than the slugs
- **3.** Hox genes are genes that control ______; we discussed how Hox mutations can set the stage for ______.
 - A. immune responses; coevolution
 - B. immune responses; adaptive radiations
 - C. body plans; coevolution
 - **D.** body plans; adaptive radiations
- **4.** Polyploidy events, where a new organism has twice as many chromosomes as the parents, are usually ______, but sometimes _____.
 - A. beneficial; lead to speciation events
 - **B.** beneficial; lead to extinction events
 - C. deleterious; create more efficient offspring
 - **D.** deleterious; create opportunities for new beneficial mutations
- **5.** A researcher wants to test the effects of supplementing flies' diets with vitamin E. They raise groups of flies in 8 different dishes. 4 of the dishes have standard diets, and 4 have supplemented diets. There are 10 flies in each dish. In this experiment, each treatment (standard and supplemented) has how many *replicates*?
 - **A.** 1
 - **B.** 4
 - **C.** 10
 - **D.** 40

Bio 1M03 Version Q

6. 80% of a frog population is destroyed by a fungal outbreak after a mild winter. What else do you need to know to determine if this was a selection event?

- **A.** Nothing. It was definitely a selection event.
- **B.** Nothing. It was definitely *not* a selection event.
- C. Whether on average the frogs that survived were in better condition (e.g., fatter or stronger) than those that died
- **D.** Whether on average the frogs that survived were different genetically than those that died
- 7. A spring storm kills almost all of the chipmunks in Hamilton. After 10 years it is found that the population is back to the original level, but has several deleterious alleles present at high frequency that were rare before the storm. This is an example of:
 - A. gene flow
 - B. speciation
 - C. a bottleneck
 - **D.** natural selection
 - E. assortative mating
- 8. Hawthorn flies have genes that attract them to either apple or hawthorn plants, and also attract them to other flies that smell like they were raised on those plants. If a population of flies lives in a place with similar amount of hawthorn and apple resources, we would expect these genotype frequencies in these attraction genes to:
 - A. Be in Hardy-Weinberg equilibrium
 - B. Be close to, but not in, Hardy-Weinberg equilibrium
 - C. Have more homozygotes than predicted by HW
 - **D.** Have more heterozygotes than predicted by HW
- **9.** Scientists studying hominin skeletons from different taxa are particularly interested in evidence of ______ but it likely reflects _____.
 - A. less robust jaws; larger brains
 - **B.** slow development; larger brains
 - C. less robust jaws; a long learning period
 - **D.** slow development; a long learning period
- 10. Two related species of plant produce flowers at different times. This likely represents
 - A. gene flow
 - **B.** balancing selection
 - C. reinforcement
 - **D.** hybridization
 - E. fusion

Student # ______

Version Q

Bio 1M03

© 2023, McMaster University Bio 1M teaching team. May be reproduced and distributed, with this notice, for non-commercial purposes only.